

Helico-conical optical beams self-heal

N. Hermosa,^{1,*} C. Rosales-Guzmán,¹ and J.P. Torres^{1,2}

¹*ICFO - Institut de Ciències Fotoniques, UPC, Mediterranean Technology Park
08860 Castelldefels (Barcelona), Spain*

²*Department of Signal Theory and Communications, UPC, 08034 Barcelona, Spain*

*Corresponding author: nathaniel.hermosa@icfo.es

Compiled November 27, 2012

An optical beam is said to be self-healing when, distorted by an obstacle, the beam corrects itself upon propagation. In this letter, we show through experiments supported by numerical simulations, that Helico-conical optical beams (HCOBs) self-heal. We observe the strong resilience of these beams with different types of obstructions, and relate this to the characteristics of their transverse energy flow. © 2012 Optical Society of America

OCIS codes: 140.3300, 260.6042, 260.0260, 070.2580.

The study of the self-healing properties of beams is of great interest in optics [1–6]. An optical beam is said to be self-healing when, after propagation, its transverse intensity profile is hardly affected by a small perturbation - a block - has been placed in its path [1–5]. The surge of interest in self-healing beams is buoyed mainly by its range of applications; self-healing can be advantageous, for instance, in beam propagation through scattering and turbulent media, and in optical manipulation [8,9].

Optical beams that exhibit self-healing include Bessel beams (BBs) [6–8], caustic beams [4], Airy beams [1,9], Pearcey beams [2], the non-paraxial Mathieu and Weber accelerating beams [10], and some forms of Laguerre-Gaussian (LG) beams [3]. In the case of the BBs and Airy beams, self-healing happens at a relatively small propagation distance, while LG beams self-heal at a distance of the order of the Rayleigh length [2,3]. Self-healing is independent of the diffracting nature of the beams, as shown by caustic [4] and LG beams [3].

In this Letter, we present another set of beams that self-heal: the Helico-conical optical beams (HCOBs). The main difference between these beams and other self-healing beams is the non-separability of their radial and azimuthal phases [11]. HCOBs possess a phase ψ that is the product of a helical phase and a conical phase: $\psi(r, \theta) = \ell\theta(K - r/r_0)$, where ℓ is the winding number around the azimuth angle θ , r_0 normalizes the radial coordinate r , and K takes either the value 0 or 1. At the far-field, the intensity profile of these beams resembles a spiral, with $K = 1$ HCOBs having a more pronounced head near the center of the beam axis compared with the $K = 0$ HCOBs. Recently, a $K = 0$ HCOB was reported to cause a spiral motion to a particle along its path [12], a three dimensional motion that combines phase gradient with intensity gradient forces [13].

It could be argued that since HCOBs have conical phases, they should behave similar to BBs. In fact, the HCOBs are more likely to be compared with fractional higher-order BBs, because of their similar intensity distributions [14]. Joint to this is the fact that HCOBs consist of strings of optical vortices upon propagation [15].




Fig. 1. (a) The experimental setup. Holograms are encoded onto a SLM. (b) Samples of unblocked (above) and blocked (below) holograms are shown. L1 and L2 are collimating lenses while M1 and M2 are mirrors for alignment. BS is beam splitter.

However, the far-field intensity pattern of experimentally generated Bessel beams, or any superposition of it, resembles a circle or a δ -ring [16], while HCOBs are spirals in the far field, and not rings when compared to BBs [11]. An important question then arises: *Can HCOBs self-heal?*

Here, we provide evidence that an HCOB reconstruct its intensity profile at a relatively short propagation distance after a small perturbation is placed in its path. We observe how the beam reconstructs for different values of ℓ and for different block sizes. Since the phase of the HCOB is not rotationally symmetric, we also note how the beam reconstructs when we change the orientation of the obstructing block. We then compare our experimental results with numerical simulations. Finally, we look at the transverse energy flow of the beam and relate it to its self-healing property.

We generate the HCOBs using a spatial light modulator (SLM) [17,18]. Figure 1(a) shows the experimental set-up. A collimated HeNe laser ($\lambda = 632.8\text{nm}$) beam impinges onto an SLM (Hamamatsu LCOS-SLM) encoded with a computer-generated hologram. The holograms ($r_0 = 2.5\text{ mm}$) are calculated from the phase of the HCOBs. A carrier frequency is added to separate the beam of interest. We observe the beam after propagation from the SLM while varying ℓ , and the size and orienta-




Fig. 2. Intensity profiles (false color, color online) of $\ell = 50$ HCOBs where (a) and (b) are obtained from experiments while (c) and (d) are from numerical simulations. The top images are for $K = 0$ while the bottom ones are for $K = 1$. The block size used in (b) and (d) is $\Delta\theta = \pi/3$.

tion of the block. We imitate the presence of a block by means of an incomplete hologram (see Fig. 1(b)). This is done for better control of the size of the block. The block size corresponds to an angular fraction $\Delta\theta$. We then capture the intensity pattern with a charged couple device (CCD) camera attached to a computer. Using the Angular Spectrum Method [20], we compare our results with numerical simulations. Figure 2 shows the intensity profiles of the unblocked and blocked HCOBs with $\ell = 50$ and $\Delta\theta = \pi/3$. The intensity patterns, not measured in the far-field, scale with the value of ℓ , similarly to what is shown in [11].

The similarities between blocked and unblocked beams are quantified for both experiments and simulations using 2D image correlation [21]. The value of the correlation coefficient ranges from 0 for non-identical beams to 1 for identical beams. We emphasize, however, that the correlation coefficient only gives a trend. It is *not* an exact measure of the quality of the beam reconstruction, especially in our case, wherein it is difficult to separate the generated beam from the adjacent diffraction orders.

The correlation coefficient changes with the value of ℓ as shown in Fig. 3(a). The block size is $\pi/3$ and the orientation of the block is shown as an inset. As the ℓ value increases, the correlation coefficient also increases in both the experiment and numerical simulations, with the $K = 0$ HCOBs having higher correlation coefficients than the $K = 1$ HCOBs with the same ℓ , at the same propagation distance.

The block size affects the reconstruction of the HCOBs as shown in Fig. 3(b). The $K = 0$ HCOBs reconstruct faster than the $K = 1$ HCOBs, given the same block size and the same propagation distance. We notice that the beam with a π block size (not shown) gives a very low correlation coefficient which is consistent with the results previously reported for other self-healing beams [4, 6]. The HCOBs reconstruction also depends on the orientation of the block as shown in Fig. 3(c). The block size is $\Delta\theta = \pi/4$ and the orientation of the block is shown below the plot. This is expected since the intensity and the phase of the beam are not rotationally symmetric.




Fig. 3. (color online) (a) Correlation for different values of ℓ . (b) Correlation with different block sizes. (c) Correlation for different block orientations. Numerical simulation (open points) uses the axis on the left, while the right axis is for experiments (filled points). Circle are for $K = 0$, and square for $K = 1$.

In another experiment, we block the HCOBs with a 0.38 mm opaque strip, and observe how the HCOBs reconstruct upon propagation. We restrict our measurement to distances below $r_0^2/\ell\lambda$, since we notice that beyond this distance, the HCOBs' intensity profiles change more rapidly [19]. Figure 4 shows the experimental and numerical results, with an opaque strip placed 16 cm after the SLM. Placing the CCD at different distances, we observe that the HCOBs self-heal as the beams propagate. The shadow of the block moves in a rotatory manner, similar to the self-healing of higher order BBs. However, unlike BBs, HCOBs slightly rotate and expand.

Figure 5 shows the transverse energy flow of an $\ell = 30$ $K = 0$ HCOB, calculated numerically [22]. Similar images can be obtained for $K = 1$ HCOBs. The direction of the energy flow traces a curved path (shown as white arrows in the figure). Even with the block, the direction




Fig. 4. (false color, color online) A 0.38mm strip is placed at the path of a $\ell = 40$ HCOB, 16cm after the SLM. (a) and (c) are experimental results while (b) and (d) are numerical simulations.




Fig. 5. (false color, color online) Transverse energy flow for $K = 0$ HCOBs with $\ell = 30$ for (a) no block, and (b) blocked located 16cm after the SLM. Both beams propagate a total distance of 20cm . Dashed lines denote the original position of the block.

of the energy flow is unaltered and so, the energy flows from the surrounding areas to the blocked area [23]. Since the energy and the energy flow are greater at the upper section, the beam reconstructs faster in this part. In addition, the transverse energy flow is greater for larger ℓ values which translates to faster reconstruction. This suggests that the transverse energy flow is the reason that the beam reconstructs.

In summary, we have shown experimentally that Helico-conical optical beams self-heal and we have supported our results with numerical simulations. The intensity profile reconstructs under different circumstances: by varying the size and the orientation of the obstructing block, as well as by changing the ℓ value of the HCOBs. We observe how the beams heal as they propagate, and we link our results with the transverse energy flow within the beam.

This work was supported by the Government of Spain (FIS2010-14831), PHORBITECH (grant number: 255914) and Fundacio Privada Cellex Barcelona. The authors would like to thank V. Rodriguez-Fajardo and L. J. Salazar-Serrano for useful discussions.

References

1. J. Broky, G. Siviloglou, A. Dogariu, and D. Christodoulides, *Opt. Express* **16**, 12880-12891 (2008). (document)
2. J. Ring, J. Lindberg, A. Mourka, M. Mazilu, K. Dholakia, and M. Dennis, *Opt. Express* **20**, 18955-18966 (2012). (document)
3. P. Vaity and R. P. Singh, *Opt. Lett.* **36**, 2994-2996 (2011). (document)
4. M. Anguiano-Morales, A. Martínez, M. Iturbe-Castillo, S. Chávez-Cerda, and N. Alcalá-Ochoa, *Appl. Opt.* **46**, 8284-8290 (2007). (document)
5. Z. Bouchal, *Opt. Comm.* **210**, 155-164 (2002). (document)
6. S. Vyas, Y. Kozawa, and S. Sata, *J. Opt. Soc. Am. A* **28**, 837-843 (2011). (document)
7. J. Durnin, *J. Opt. Soc. Am. A* **4**, 651-654 (1987). (document)
8. V. Garcés-Chávez, D. McGloin, H. Melville, W. Sibbett, and K. Dholakia, *Nature* **419**, 145-147 (2002). (document)
9. J. Baumgartl, M. Mazilu, and K. Dholakia, *Nature Photonics* **2**, 675 - 678 (2008). (document)
10. P. Zhang, Y. Hu, T. Li, D. Cannan, X. Yin, R. Morandotti, Z. Chen, X. Zhang, *Phys. Rev. Lett.* **109**, 193901 (2012). (document)
11. C. Alonzo, P. Rodrigo, and J. Glückstad, *Opt. Express* **13**, 1749-1760 (2005). (document)
12. V. R. Daria, D. Z. Palima, and J. Glückstad, *Opt. Express* **19**, 476-481 (2011). (document)
13. M. Padgett, and R. Bowman, *Nature Photonics* **5**, 343-347 (2011). (document)
14. S.H. Tao, W.M. Lee, and X. Yuan, *Appl. Opt.* **43**, 122-126 (2004). (document)
15. N. Hermosa, and C.O. Manaiois, *Opt. Comm.* **271**, 178-183 (2007). (document)
16. L.C. Thomson, and J. Courtial, *Opt. Comm.* **281**, 1217-1221 (2007). (document)
17. A. Vasara, J. Turunen, and A. Friberg, *J. Opt. Soc. Am. A* **6**, 1748-1754 (1989). (document), 19, 19
18. N. Chattrapiban, E. Rogers, D. Cofield, W. Hill, III, and R. Roy, *Opt. Lett.* **28**, 2183-2185 (2003). (document)
19. This formula comes from the maximum propagation distance of a BB [17]. For HCOBs, the slope of the cone depends on ℓ , and the smallest value of the propagation distance is at it's steepest slope. The steepest slope thus define the propagation distance of the HCOBs. (document)
20. J.W. Goodman, *Introduction to Fourier Optics 3rd ed.* (Roberts & Company, 2005). (document)
21. N. Matsumoto, T. Ando, T. Inoue, Y. Ohtake, N. Fukuchi, and T. Hara, *J. Opt. Soc. Am. A* **25**, 1642-1651 (2008). (document)
22. L. Allen, M.W. Beijersbergen, R.J.C. Spreeuw, and J.P. Woerdman, *Phys. Rev. A* **45**, 8185-8189 (1992). (document)
23. A. Bekshaev, and M. Soskin, L. Allen, *Opt. Lett.* **31**, 2100-2201 (2006). (document)

Informational Fourth Page

In this section, please provide full versions of citations to assist reviewers and editors (OL publishes a short form of citations) or any other information that would aid the peer-review process.

References

1. J. Broky, G. Siviloglou, A. Dogariu, and D. Christodoulides, "Self-healing properties of optical Airy beams," *Opt. Express* **16**, 12880-12891 (2008).
2. J. Ring, J. Lindberg, A. Mourka, M. Mazilu, K. Dholakia, and M. Dennis, "Auto-focusing and self-healing of Pearcey beams," *Opt. Express* **20**, 18955-18966 (2012).
3. P. Vaity and R. P. Singh, "Self-healing property of optical ring lattice," *Opt. Lett.* **36**, 2994-2996 (2011).
4. M. Anguiano-Morales, A. Martínez, S. Chávez-Cerda, and N. Alcalá-Ochoa, "Self-healing property of a caustic optical beam," *Appl. Opt.* **46**, 8284-8290 (2007).
5. Z. Bouchal, "Resistance of nondiffracting vortex beam against amplitude and phase perturbations," *Opt. Comm.* **210**, 155-164 (2002).
6. S. Vyas, Y. Kozawa, and S. Sata, "Self-healing of tightly focused scalar and vector Bessel-Gauss beams at the focal plane," *J. Opt. Soc. Am. A* **28**, 837-843 (2011).
7. J. Durnin, "Exact solutions for nondiffracting beams. I. The scalar theory," *J. Opt. Soc. Am. A* **4**, 651-654 (1987).
8. V. Garcés-Chávez, D. McGloin, H. Melville, W. Sibbett, and K. Dholakia, "Simultaneous micromanipulation in multiple planes using a self-reconstructing light beam," *Nature* **419**, 145-147 (2002).
9. J. Baumgartl, M. Mazilu, and K. Dholakia, "Optically mediated particle clearing using Airy wavepackets," *Nature Photonics* **2**, 675 - 678 (2008).
10. P. Zhang, Y. Hu, T. Li, D. Cannan, X. Yin, R. Morandotti, Z. Chen, X. Zhang, "Nonparaxial Mathieu and Weber accelerating beams," *Phys. Rev. Lett.* **109**, 193901 (2012).
11. C. Alonzo, P. Rodrigo, and J. Glückstad, "Helico-conical optical beams: a product of helical and conical phase fronts," *Opt. Express* **13**, 1749-1760 (2005).
12. V. R. Daria, D. Z. Palima, and J. Glückstad, "Optical twists in phase and amplitude," *Opt. Express* **19**, 476-481 (2011).
13. M. Padgett, and R. Bowman, "Tweezers with a twist," *Nature Photonics* **5**, 343-347 (2011).
14. S.H. Tao, W.M. Lee, and X. Yuan, "Experimental study of holographic generation of fractional Bessel beams," *Appl. Opt.* **43**, 122-126 (2004).
15. N. Hermosa, and C.O. Manaois, "Phase structure of helico-conical optical beams," *Opt. Comm.* **271**, 178-183 (2007).
16. L.C. Thomson, and J. Courtial, "Holographic shaping of generalized self-reconstructing light beams," *Opt. Comm.* **281**, 1217-1221 (2007).
17. A. Vasara, J. Turunen, and A. Friberg, "Realization of general nondiffracting beams with computer-generated holograms," *J. Opt. Soc. Am. A* **6**, 1748-1754 (1989).
18. N. Chattrapiban, E. Rogers, D. Cofield, W. Hill, III, and R. Roy, "Generation of nondiffracting Bessel beams by use of a spatial light modulator," *Opt. Lett.* **28**, 2183-2185 (2003).
19. This formula comes from the maximum propagation distance of a BB [17]. For HCOBs, the slope of the cone depends on ℓ , and the smallest value of the propagation distance is at its steepest slope. The steepest slope thus define the propagation distance of the HCOBs.
20. J.W. Goodman, *Introduction to Fourier Optics 3rd ed.* (Roberts & Company, 2005).
21. N. Matsumoto, T. Ando, T. Inoue, Y. Ohtake, N. Fukuchi, and T. Hara, "Generation of high-quality higher-order Laguerre-Gaussian beams using liquid-crystal-on-silicon spatial light modulators," *J. Opt. Soc. Am. A* **25**, 1642-1651 (2008).
22. L. Allen, M.W. Beijersbergen, R.J.C. Spreeuw, and J.P. Woerdman, "Orbital angular momentum of light and the transformation of Laguerre-Gaussian laser modes," *Phys. Rev. A* **45**, 8185-8189 (1992).
23. A. Bekshaev, and M. Soskin, L. Allen, "Rotational transformations and transverse energy flow in paraxial light beams: linear azimuthons," *Opt. Lett.* **31**, 2100-2201 (2006).